



## Preventive Care/Immunization Schedule

|           |  |
|-----------|--|
| Birth     | Hepatitis B  |
| 1-2 weeks | Newborn Screening (Hep B if not given at birth)                |
| 1 month   | None   |
| 2 months  | Vaxelis (DTaP-IPV-Hib-Hep B), PCV13, Rotavirus                 |
| 4 months  | Vaxelis (DTaP-IPV-Hib-Hep B), PCV13, Rotavirus                 |
| 6 months  | Vaxelis (DTaP-IPV-Hib-Hep B), PCV13, Rotavirus, Influenza(Flu) |
| 9 months  | None   |
| 12 months | MMR, Varicella (Chickenpox), Hepatitis A                       |
| 15 months | Pentacel (DTaP-IPV-HIB), PCV13                                 |
| 18 months | None   |
| 24 months | Hepatitis A  |
| 30 months | None   |
| 3 years   | None   |
| 4 years   | Quadracel (DTaP-Polio) ProQuad (MMR-Varicella)                 |
| 5 years   | None   |
| 6 years   | None   |
| 7 years   | None   |
| 8 years   | None   |
| 9 years   | None   |
| 10 years  | None   |
| 11 years  | HPV, Tdap, MenACWY   |
| 12 years  | HPV  |
| 13 years  | None   |
| 14 years  | None   |
| 15 years  | None   |
| 16 years  | MenACWY, MenB  |
| 17 years  | MenB   |
| 18 years  | None   |

### ***Vaccine Index:***

- **Chickenpox:** 1<sup>st</sup> dose given individually, 2<sup>nd</sup> dose given as a combo vaccine with MMR, (MMR/Chickenpox)
- **DTaP/Polio:** Diphtheria, Tetanus, Pertussis, Poliomyelitis (Quadracel)
- **DTaP-Polio-Hib-Hep B:** Diphtheria, Tetanus, Pertussis, Poliomyelitis, Hepatitis B, Haemophilus Influenzae type b; series of 3-4
- **Hepatitis A:** series of 2
- **HPV:** Human Papillomavirus; series of 2 (if given before age 15)
- **MenACWY:** Meningococcal meningitis, types A, C, W, and Y; series of 2
- **MenB:** Meningococcal meningitis, type B; series of 2
- **MMR:** Measles, Mumps, Rubella (1<sup>st</sup> dose given individually, 2<sup>nd</sup> dose given as a combo vaccine with Chickenpox)
- **Pro Quad:** Measles, Mumps, Rubella, Chickenpox (given for the 2<sup>nd</sup> doses for both MMR & Chickenpox)
- **PCV13:** pneumococcal conjugate; series of 4
- **Rotavirus:** series of 3
- **Tdap:** Tetanus, Diphtheria, Pertussis (for children 7 or older)
- **Influenza:** minimum age 6months- 2 doses, separated by at least 4 weeks, for children age 6 months-8 years who have not received 2 doses in the same consecutive flu season.